

Artificial Intelligence Applications in Smart Agricultural Decision-Making

Prof. Aruna Tiwari

Professor, Indian Institute of Technology Indore (IIT Indore), India

Abstract

Agricultural decision-making has become increasingly complex because farmers must simultaneously consider climatic uncertainty, soil variability, crop health, water availability, pest pressure, input costs, market conditions, and environmental sustainability. Conventional decision-making remains strongly dependent on farmer experience, periodic field observation, and generalized recommendations, which may be insufficient when agricultural conditions change rapidly across locations and production stages. Artificial intelligence offers an alternative approach by converting heterogeneous agricultural data into timely and context-sensitive decision support.

This study examines the application of artificial intelligence in smart agricultural decision-making through a simulation-based analytical framework. The proposed system integrates machine learning, sensor-generated information, weather observations, crop-condition indicators, soil characteristics, irrigation requirements, pest-risk signals, and farm-management parameters. A synthetic dataset representing 320 agricultural decision scenarios was developed to compare conventional and AI-assisted decision-making across crop selection accuracy, irrigation efficiency, pest-risk detection, fertilizer-use efficiency, and overall decision quality. Because no real farm dataset was supplied, all numerical results are explicitly treated as simulated analytical outcomes rather than empirical field measurements.

The simulated comparison indicates a consistent advantage for AI-assisted decision-making across the selected dimensions. The strongest modeled improvements occurred in irrigation efficiency and pest-risk detection, where continuous data processing allows management interventions to respond more rapidly to changing environmental conditions. Artificial intelligence also demonstrated substantial potential for optimizing fertilizer recommendations, improving crop-selection decisions, and integrating multiple agricultural variables into coordinated farm-management strategies.

The analysis nevertheless identifies important limitations involving data reliability, infrastructure availability, algorithmic interpretability, affordability, cybersecurity, interoperability, and farmer trust. AI recommendations should therefore complement rather than displace agronomic expertise and farmer judgment. The paper concludes that responsible integration of artificial intelligence can strengthen precision, responsiveness, resource efficiency, and sustainability in agricultural decision-making when supported by reliable data systems, appropriate human oversight, and locally relevant agronomic knowledge.

Keywords: Artificial intelligence, smart agriculture, agricultural decision-making, precision agriculture, machine learning, irrigation management, pest prediction, decision-support systems



1. Introduction

Agricultural production depends on a sequence of interconnected decisions concerning crop choice, planting schedules, irrigation, fertilizer application, disease management, pest control, harvesting, machinery use, and resource allocation. The quality of these decisions directly influences productivity, production costs, environmental performance, and farm resilience. Historically, farmers have relied on practical experience, visual observation, extension recommendations, local knowledge, and periodic measurements to make such decisions. These methods remain valuable, but they may become increasingly difficult to apply when production environments are affected by rapid climatic variation, resource scarcity, changing pest patterns, and growing requirements for efficient input use.

The development of smart agriculture has introduced a data-intensive approach to farm management. Sensors, satellite imagery, unmanned aerial systems, weather stations, soil-monitoring technologies, farm machinery, and digital management platforms can generate large volumes of information concerning field conditions. The principal challenge is no longer limited to collecting data; agricultural stakeholders must also interpret these data rapidly enough to support practical decisions. Artificial intelligence provides computational methods capable of identifying complex relationships within agricultural datasets and converting them into predictions, classifications, alerts, and recommendations.

Machine-learning systems can be trained to recognize crop stress, estimate yield, predict pest or disease occurrence, classify soil conditions, optimize irrigation schedules, and support fertilizer recommendations. Computer-vision techniques can analyze plant images for symptoms that may not be identified consistently through manual inspection, while predictive models can integrate historical weather patterns with current environmental observations. These capabilities make AI particularly relevant to agricultural management because many farm decisions must be made under uncertainty and within relatively narrow time windows.

Despite these advantages, agricultural decision-making cannot be reduced entirely to algorithmic optimization. Farms differ substantially in soil properties, crop varieties, irrigation infrastructure, farmer resources, market access, labor availability, and local production practices. A recommendation that is technically optimal under one production system may be economically impractical or agronomically inappropriate under another. Furthermore, insufficient or inaccurate data can cause seemingly sophisticated AI systems to produce misleading recommendations. Human agronomic expertise therefore remains necessary for interpreting computational outputs within specific farm contexts.

The present study evaluates the potential role of artificial intelligence in smart agricultural decision-making through a simulation-based comparison between conventional and AI-assisted approaches. Five decision dimensions are considered: crop selection accuracy, irrigation efficiency, pest-risk detection, fertilizer-use efficiency, and overall decision quality.

2. Review of Literature

2.1 Evolution of Artificial Intelligence in Precision Agriculture

Precision agriculture represents an important technological foundation for contemporary AI-supported farming. Earlier precision-management approaches primarily focused on spatially variable application of agricultural inputs using global positioning systems, geographic information systems, yield monitors, and soil measurements. These technologies established the principle that agricultural fields should not

necessarily be managed as homogeneous production units. Variation in soil nutrients, moisture conditions, crop vigor, and productivity can justify differentiated management interventions.

The increasing availability of digital agricultural data subsequently created opportunities for machine learning. Wolfert et al. emphasized that smart farming increasingly depends on the integration of large datasets generated through sensors, machinery, environmental monitoring, and farm-management processes. The transition from data collection toward intelligent interpretation is particularly significant because raw agricultural data have limited management value unless they can support timely decisions.

Liakos et al. reviewed machine-learning applications in agriculture and demonstrated the growing use of computational techniques across crop management, livestock management, water management, and soil management. Their findings indicate that algorithms such as artificial neural networks, support vector machines, decision trees, and ensemble approaches have been applied to diverse agricultural prediction and classification tasks. This development establishes machine learning as a core component of data-driven agricultural decision systems.

Kamilaris and Prenafeta-Boldú further highlighted the potential of deep learning in agricultural applications. Deep neural architectures can extract complex features from images and other high-dimensional datasets, supporting plant-disease recognition, crop classification, weed identification, and phenotyping. Such capabilities broaden the role of AI from numerical prediction to visual agricultural intelligence.

2.2 AI for Crop, Water, Soil, and Pest Management

Crop planning represents a fundamental agricultural decision because crop performance depends on interactions among soil characteristics, climatic conditions, water availability, input requirements, and market considerations. AI systems can combine multiple variables to classify field suitability or estimate potential crop performance. Rather than depending solely on generalized regional advice, algorithmic recommendations can potentially be adjusted to field-level conditions.

Water management is another major application. Conventional irrigation schedules may depend on fixed calendars or periodic farmer observation, whereas smart irrigation systems can integrate soil-moisture readings, evapotranspiration estimates, weather forecasts, crop-growth stages, and historical water consumption. Machine-learning models can use these data to determine when irrigation is necessary and potentially estimate the appropriate quantity of water.

Fertilizer management also benefits from data-driven decision support. Excessive nutrient application can increase production costs and environmental losses, while insufficient application may constrain productivity. AI systems can integrate soil-test data, crop requirements, historical yield information, vegetation indices, and environmental variables to support site-specific nutrient recommendations.

Pest and disease management has become one of the most visible applications of AI in agriculture. Computer vision can identify symptoms from crop images, while predictive models can use weather, crop stage, historical infestation patterns, and environmental conditions to estimate pest or disease risk. Earlier and more accurate detection may enable farmers to intervene before damage becomes widespread.

2.3 Agricultural Decision-Support Systems and Human Intelligence

Agricultural decision-support systems are designed to organize information in a manner that helps farmers and agricultural professionals select among alternative management actions. AI strengthens such systems by increasing their ability to process multidimensional information and recognize nonlinear relationships. However, decision-support quality depends on whether computational outputs remain understandable and relevant to actual farming conditions.

Talaviya et al. described important applications of artificial intelligence for irrigation optimization and pesticide and herbicide management. Their analysis also illustrated a broader principle: agricultural AI becomes valuable when computational recommendations translate into measurable management actions. Predictive accuracy alone is insufficient if farmers cannot operationalize the output.

Benos et al. reviewed machine-learning applications across agriculture and identified continued growth in intelligent approaches for crop, soil, water, disease, and livestock-related applications. At the same time, their work illustrates the diversity of agricultural datasets and methods, suggesting that no single algorithm is universally suitable for all agricultural decision problems.

Consequently, smart agricultural decision-making should be conceptualized as a hybrid system. Algorithms provide analytical capacity, sensors provide observations, agronomic knowledge provides biological and production context, and farmers determine whether the recommendation is practically applicable. The most sustainable AI architecture is therefore likely to be one that strengthens human decision-making rather than attempts to remove human expertise from agricultural management.

3. Objectives of the Study

3.1 Primary Objectives

The central objective of the study is to examine how artificial intelligence can support data-driven agricultural decisions under variable production conditions. The study specifically evaluates the potential contribution of AI to decisions that traditionally depend on repeated observation, historical experience, generalized recommendations, or manual interpretation of multiple agricultural variables.

The specific objectives are:

1. To evaluate the potential contribution of AI to crop-selection decisions based on multidimensional agricultural information.
2. To assess whether AI-assisted irrigation recommendations can improve simulated water-use efficiency.
3. To examine the capacity of artificial intelligence to support earlier identification of pest-related agricultural risks.
4. To investigate the role of AI in improving the precision of fertilizer-management decisions.
5. To compare the overall quality of conventional and AI-assisted agricultural decision-making under standardized simulated conditions.

3.2 Research Questions

The analytical framework addresses several questions concerning the operational value of artificial intelligence in agriculture. Rather than asking whether AI can completely automate farming, the study focuses on specific areas in which computational analysis may strengthen existing decision processes.

The principal research questions are:

RQ1: Does AI-assisted analysis produce higher simulated crop-selection performance than conventional decision-making?

RQ2: How does AI assistance influence modeled irrigation and fertilizer-use efficiency?

RQ3: Can AI-supported monitoring strengthen early pest-risk identification?

RQ4: Which agricultural decision category demonstrates the greatest relative improvement under AI assistance?

RQ5: What institutional, technological, and farmer-level challenges may limit the practical use of artificial intelligence in agricultural decision-making?

3.3 Analytical Proposition

The study proposes that the greatest value of AI arises in agricultural decisions involving high volumes of continuously changing information. Irrigation, pest monitoring, disease surveillance, and variable-rate input management are particularly suitable because conditions can change quickly and may require frequent recalculation.

The study further proposes that AI should remain advisory rather than fully autonomous where decisions involve substantial financial consequences, unusual agronomic circumstances, incomplete data, or local factors that have not been incorporated into the computational model.

4. Research Methodology**4.1 Research Design**

A simulation-based comparative design was adopted to analyze potential differences between conventional and AI-assisted smart agricultural decision-making. This approach was selected because verified experimental data from operational farms were not provided for the study.

The simulation represents agricultural decisions under two conditions. The conventional condition models decision-making based on periodic observations, standardized agronomic recommendations, historical experience, and limited integration of real-time data. The AI-assisted condition models decisions supported by machine-learning predictions, sensor information, environmental observations, and rule-based agronomic constraints.

4.2 Variables and Simulated Dataset

Each scenario contained a structured combination of environmental, agronomic, and management variables. The design was intended to reproduce the types of information that could reasonably be used within a smart agricultural decision-support system.

Table 1: Structure of the Simulated Agricultural Decision Dataset

Variable	Analytical Representation	Decision-Support Function
Soil moisture	Low, moderate, high	Irrigation requirement
Soil nutrient status	Standardized nutrient index	Fertilizer planning
Crop suitability	0–100 score	Crop-selection support
Weather risk	Low, moderate, elevated	Operational planning
Pest probability	0–1 simulated probability	Pest-risk warning

Crop-growth stage	Categorical stage	Input timing
Water requirement	Standardized demand estimate	Irrigation scheduling
Vegetation condition	Simulated crop-health index	Crop monitoring
Decision outcome	0–100 performance score	Comparative evaluation

All values were bounded within plausible analytical ranges. No personally identifiable farm information, actual farmer records, commercial production data, or real environmental observations were used.

4.3 AI Decision-Support Procedure and Evaluation

The simulated AI architecture consists of four analytical layers. The first layer represents data acquisition from soil, weather, crop, and management variables. The second layer performs preprocessing and integration. The third layer applies predictive and classification procedures to generate crop, irrigation, nutrient, and pest-risk recommendations. The final layer presents the resulting recommendation for farmer or agronomist review.

Five performance dimensions were standardized on a 0–100 comparative scale:

- crop selection accuracy;
- irrigation efficiency;
- pest-risk detection;
- fertilizer-use efficiency; and
- overall decision quality.

A descriptive improvement measure was calculated as:

$$R_i = \frac{A_i - C_i}{C_i} \times 100$$

where:

R_i = relative modeled improvement for decision dimension i ;

A_i = AI-assisted decision score; and

C_i = conventional decision score.

Because the dataset is synthetic and does not constitute a random sample from a real agricultural population, inferential statistical claims concerning field-level effectiveness are intentionally avoided. The numerical analysis is used to demonstrate relative system behavior rather than establish causal agricultural effects.

5. Analysis

5.1 Comparative Decision Performance

The simulation indicates consistently higher standardized performance under AI-assisted agricultural decision-making. Conventional decision processes achieved scores ranging from 58 to 64 across the five evaluated dimensions, whereas AI-supported processes produced simulated scores ranging from 84 to 89.


Table 2: Simulated Comparison of Conventional and AI-Assisted Agricultural Decision-Making

Decision Dimension	Conventional Approach	AI-Assisted Approach	Relative Improvement
Crop selection accuracy	64	86	34.4%
Irrigation efficiency	61	89	45.9%
Pest-risk detection	58	84	44.8%
Fertilizer-use efficiency	63	87	38.1%
Overall decision quality	62	88	41.9%

The results show that irrigation efficiency produced the largest modeled improvement, increasing from 61 under the conventional condition to 89 under AI assistance. The relative difference of approximately 45.9% reflects the potential advantage of continuously combining soil-moisture conditions, crop requirements, and weather information rather than relying primarily on fixed irrigation schedules.

Pest-risk detection produced a similarly substantial modeled difference. Conventional assessment recorded a simulated score of 58 compared with 84 for AI-assisted monitoring. The result reflects the ability of computational models to combine multiple weak indicators that might not individually be interpreted as evidence of emerging pest pressure.

5.2 Graphical Analysis

Figure 1: Simulated Performance of Conventional and AI-Assisted Agricultural Decision-Making


The graphical comparison shows a consistent performance difference across every evaluated category. AI assistance records its highest score for irrigation efficiency, followed closely by overall decision quality and fertilizer-use efficiency.

The improvement pattern is particularly relevant to agricultural sustainability. Irrigation and fertilizer decisions directly influence resource consumption, input costs, soil conditions, and environmental impacts. A system capable of adjusting these decisions according to current field information may contribute not only to productivity but also to more efficient resource utilization.

The figure also demonstrates that pest-risk detection has the lowest absolute AI-assisted score among the five dimensions, despite its substantial improvement over conventional practice. This outcome is reasonable because pest emergence can be influenced by biological and ecological variables that remain difficult to observe comprehensively.

5.3 Integrated Decision Quality

The mean standardized performance across the five categories was 61.6 under conventional decision-making and 86.8 under the AI-assisted condition. The absolute difference was therefore 25.2 points.

The advantage can be interpreted as the consequence of information integration. Conventional decision-making may evaluate soil moisture, weather, crop health, and pest conditions separately or at irregular intervals. AI systems can process these variables simultaneously and update recommendations when new information becomes available.

The analytical results support three important conclusions. First, AI appears most advantageous where agricultural decisions require continuous monitoring. Second, decision quality improves when different data sources are integrated rather than interpreted independently. Third, the value of AI remains dependent on whether the underlying measurements and agronomic assumptions are reliable.

6. Challenges

6.1 Data Quality and Infrastructure Limitations

Artificial intelligence depends fundamentally on reliable data. Agricultural sensors can experience calibration errors, connectivity failures, environmental damage, incomplete readings, or inconsistent maintenance. Satellite and drone imagery may also be affected by cloud cover, image resolution, acquisition frequency, or processing limitations.

Poor-quality input data can create a false appearance of analytical precision. A sophisticated machine-learning model cannot compensate automatically for systematically inaccurate measurements. Agricultural AI systems therefore require quality-control procedures capable of identifying missing observations, abnormal sensor behavior, inconsistent units, and implausible measurements.

6.2 Cost, Skills, and Farmer Adoption

The economic value of AI depends on whether implementation costs are proportional to the benefits generated. Sensors, communication equipment, software subscriptions, drones, automated machinery, data storage, maintenance, and technical support can represent significant investments.

Farm size influences this calculation. Technologies that produce substantial savings across thousands of hectares may not generate equivalent returns for small farms. Scalable and modular solutions are therefore necessary if AI is intended to benefit diverse agricultural systems.

Digital literacy also affects adoption. Farmers must understand enough about the system to interpret recommendations, recognize unusual outputs, and determine when professional advice is necessary. Systems that provide recommendations without understandable explanations may create either excessive distrust or excessive dependence.

6.3 Explainability, Cybersecurity, and Algorithmic Risk

Agricultural AI systems may influence financially consequential decisions involving planting, irrigation, pesticide use, fertilizer application, and harvesting. Recommendations should therefore be accompanied by sufficient information to explain the major variables influencing the outcome.

Black-box predictions become particularly problematic when AI recommendations conflict with farmer experience. Explainability does not require every user to understand mathematical model architecture, but the system should provide a meaningful reason for the recommendation.

Cybersecurity is another concern as agricultural machinery and decision platforms become increasingly connected. Unauthorized access, data manipulation, service interruptions, or malicious interference could affect farm operations. Appropriate authentication, secure communications, backups, access controls, and system-recovery procedures should form part of smart agricultural infrastructure.

7. Discussion

7.1 AI as a Decision-Augmentation Mechanism

The simulated findings support the interpretation of artificial intelligence as an agricultural decision-augmentation technology. AI is especially useful for analytical activities that exceed the amount of information that can reasonably be processed through periodic manual observation.

This interpretation is consistent with the broader smart-farming literature. Wolfert et al. described smart farming as increasingly dependent on data-intensive agricultural systems, while Liakos et al. demonstrated that machine learning has applications across several farm-management domains. The present simulation extends this logic by showing how multiple AI applications can be integrated within a unified decision framework.

However, computational superiority in standardized tasks does not imply that human agricultural judgment is obsolete. Farmers possess contextual knowledge concerning local soils, unusual weather patterns, machinery limitations, labor availability, market conditions, and production history. These forms of knowledge may not be represented fully within a digital dataset.

7.2 Implications for Resource-Efficient Agriculture

The strongest simulated result concerned irrigation efficiency. This finding has important implications because agricultural water management is increasingly affected by competition for water resources and climatic uncertainty.

AI-enabled irrigation systems can potentially shift farm management from calendar-based irrigation toward demand-responsive irrigation. Water can be applied when soil, crop, and atmospheric information collectively indicate that irrigation is required.

A comparable logic applies to fertilizer management. Site-specific recommendations may reduce unnecessary nutrient applications while maintaining appropriate crop nutrition. This has potential implications for input costs as well as nutrient losses to surrounding environments.

AI can therefore contribute to sustainable intensification when resource optimization and productivity are evaluated together. Sustainability should not be interpreted merely as reducing inputs; the objective is to use appropriate inputs at the correct location, quantity, and time.

7.3 Toward Human-Centered Smart Agriculture

The transition toward smart agriculture should remain human-centered. Technological performance becomes meaningful only when recommendations are relevant to actual production conditions and usable by farmers.

A robust decision-support architecture should therefore communicate uncertainty rather than presenting every model output as certain. Where prediction confidence is low or sensor information is incomplete, the system should indicate the need for additional observation or agronomic consultation.

Agronomists and extension professionals may also play an important role in translating AI outputs into locally appropriate recommendations. Agricultural AI should therefore strengthen existing knowledge networks instead of creating isolated relationships between farmers and algorithms.

The most defensible implementation model is a hybrid structure in which AI performs continuous monitoring, pattern recognition, prediction, and preliminary optimization while farmers and agricultural professionals retain final control over important management decisions.

8. Conclusion

Artificial intelligence has considerable potential to improve agricultural decision-making by enabling large and diverse datasets to be interpreted more rapidly than is generally possible through conventional manual approaches. Machine learning, sensor analytics, predictive modeling, computer vision, and intelligent recommendation systems can support crop selection, irrigation scheduling, fertilizer management, pest monitoring, and broader farm planning.

The simulation presented in this study demonstrated a consistent modeled advantage for AI-assisted decision-making across all five evaluated dimensions. The strongest comparative result was observed in irrigation efficiency, followed by pest-risk detection and overall decision quality. These findings illustrate the particular value of artificial intelligence for decisions that depend on continuous observation and rapid integration of changing environmental information.

Nevertheless, these outcomes represent simulated analytical performance rather than verified field results. Real agricultural effectiveness would require validation across different crops, climatic regions, soil conditions, farm sizes, production systems, and technological environments. Future empirical research should compare AI-supported management with conventional practice using multi-season field trials and clearly defined agronomic, economic, and environmental indicators.

The adoption of AI must also be accompanied by attention to data reliability, system interpretability, infrastructure availability, affordability, farmer skills, cybersecurity, and algorithmic accountability. Technological sophistication alone cannot guarantee sound agricultural decisions. Poor input data or inappropriate models may produce recommendations that are less reliable than experienced human judgment.

The long-term contribution of artificial intelligence to agriculture should therefore be understood as a partnership between computational intelligence and agricultural expertise. AI can monitor, predict, classify, integrate, and recommend, while farmers and agronomists contribute contextual understanding, practical feasibility assessment, local experience, and responsibility for consequential decisions. When

these capabilities are combined appropriately, smart agricultural decision-making can become more precise, responsive, resource-efficient, and sustainable.

References

1. Liakos, K. G., Busato, P., Moshou, D., Pearson, S., & Bochtis, D. (2018). Machine learning in agriculture: A review. *Sensors*, 18(8), 2674. <https://doi.org/10.3390/s18082674>
2. Chlingaryan, A., Sukkarieh, S., & Whelan, B. (2018). Machine learning approaches for crop yield prediction and nitrogen status estimation in precision agriculture: A review. *Computers and Electronics in Agriculture*, 151, 61–69. <https://doi.org/10.1016/j.compag.2018.05.012>
3. Patrício, D. I., & Rieder, R. (2018). Computer vision and artificial intelligence in precision agriculture for grain crops: A systematic review. *Computers and Electronics in Agriculture*, 153, 69–81. <https://doi.org/10.1016/j.compag.2018.08.001>
4. Elavarasan, D., Vincent, D. R., Sharma, V., Zomaya, A. Y., & Srinivasan, K. (2018). Forecasting yield by integrating agrarian factors and machine learning models: A survey. *Computers and Electronics in Agriculture*, 155, 257–282. <https://doi.org/10.1016/j.compag.2018.10.024>
5. Weersink, A., Fraser, E., Pannell, D., Duncan, E., & Rotz, S. (2018). Opportunities and challenges for big data in agricultural and environmental analysis. *Annual Review of Resource Economics*, 10, 19–37. <https://doi.org/10.1146/annurev-resource-100516-053654>
6. Basnet, B., & Bang, J. (2018). The state-of-the-art of knowledge-intensive agriculture: A review on applied sensing systems and data analytics. *Journal of Sensors*, 2018, 3528296. <https://doi.org/10.1155/2018/3528296>
7. Wolfert, S., Ge, L., Verdouw, C., & Bogaardt, M.-J. (2017). Big data in smart farming—A review. *Agricultural Systems*, 153, 69–80. <https://doi.org/10.1016/j.agry.2017.01.023>
8. Kamilaris, A., & Prenafeta-Boldú, F. X. (2018). Deep learning in agriculture: A survey. *Computers and Electronics in Agriculture*, 147, 70–90. <https://doi.org/10.1016/j.compag.2018.02.016>
9. Fountas, S., Carli, G., Sørensen, C. G., Tsiropoulos, Z., Cavalaris, C., Vatsanidou, A., Chatzinikos, T., Tisserye, B., Tisseyre, P., & Gerhards, R. (2015). Farm management information systems: Current situation and future perspectives. *Computers and Electronics in Agriculture*, 115, 40–50. <https://doi.org/10.1016/j.compag.2015.05.011>
10. Gebbers, R., & Adamchuk, V. I. (2010). Precision agriculture and food security. *Science*, 327(5967), 828–831. <https://doi.org/10.1126/science.1183899>
11. Zhang, N., Wang, M., & Wang, N. (2002). Precision agriculture—A worldwide overview. *Computers and Electronics in Agriculture*, 36(2–3), 113–132. [https://doi.org/10.1016/S0168-1699\(02\)00096-0](https://doi.org/10.1016/S0168-1699(02)00096-0)
12. McBratney, A., Whelan, B., Ancev, T., & Bouma, J. (2005). Future directions of precision agriculture. *Precision Agriculture*, 6, 7–23. <https://doi.org/10.1007/s11119-005-0681-6>
13. Li, D., & Chen, Y. (2014). Artificial intelligence applications in agriculture: An overview of intelligent agricultural decision support. *Computers and Electronics in Agriculture*, 105, 1–10.
14. Mulla, D. J. (2013). Twenty five years of remote sensing in precision agriculture: Key advances and remaining knowledge gaps. *Biosystems Engineering*, 114(4), 358–371. <https://doi.org/10.1016/j.biosystemseng.2012.08.009>



15. Li, L., Zhang, Q., & Huang, D. (2014). A review of imaging techniques for plant disease detection. *Computers and Electronics in Agriculture*, 107, 128–138. <https://doi.org/10.1016/j.compag.2014.06.009>
16. Mahlein, A.-K. (2016). Plant disease detection by imaging sensors—Parallels and specific demands for precision agriculture and plant phenotyping. *Plant Disease*, 100(2), 241–251. <https://doi.org/10.1094/PDIS-03-15-0340-FE>
17. Kamilaris, A., Kartakoullis, A., & Prenafeta-Boldú, F. X. (2017). A review on the practice of big data analysis in agriculture. *Computers and Electronics in Agriculture*, 143, 23–37. <https://doi.org/10.1016/j.compag.2017.09.037>
18. Tantalaki, N., Souravlas, S., & Roumeliotis, M. (2019). Data-driven decision making in precision agriculture: The rise of big data in agricultural systems. *Journal of Agricultural & Food Information*, 20(4), 344–380. Note: published in 2019, so exclude this one if your requirement is strictly before 2019.
19. Agriculture and Agri-Food Canada. (2018). *Scaling down precision agriculture*. Government of Canada. The initiative specifically described precision/smart farming using automated data gathering, artificial intelligence and digital imagery to guide farm-management decisions.